

Upper Columbia Steelhead ESU

Artificial Propagation Review

Kristine Petersen

Salmon Recovery Division





Included in the ESU

- ◆ Integrated Programs
- ◆ Conservation & Harvest Mitigation
 - Wenatchee Basin
 - ◆ Wenatchee River stock \approx 400K smolts
 - Methow Basin
 - ◆ Winthrop NFH (Wells stock) \approx 100K smolts
 - ◆ WDFW (Wells stock) \approx 250K smolts
 - Okanogan Basin
 - ◆ WDFW (Wells stock) \approx 150K smolts
 - ◆ Colville Tribes \approx 40K smolts


Included in the ESU (continued)

- ◆ Other Programs
- ◆ Harvest Mitigation
 - Mainstem Columbia River
 - ◆ Ringold Hatchery (Wells stock) \approx 180K smolts

ESA Authorization

UCR steelhead artificial propagation programs are authorized under ESA Section 10 Permits 1396, 1396, and 1412 which require:

“the taking will not appreciably reduce the likelihood of the survival and recover of the species in the wild” (ESA Section 10(a)(B)(iv))

A stylized, layered mountain range graphic in shades of teal and blue, located in the bottom right corner of the slide.

“The effects of hatchery fish on the likelihood of extinction of an ESU, depend on how hatchery fish affect four key attributes”




Abundance

- ◆ Wenatchee Basin (1996-2003)
 - 1,220 natural steelhead adults
 - 1,525 hatchery steelhead adults
 - Average of 43% hatchery origin
- ◆ Above Wells Dam (1986–2003)
 - 455 natural steelhead adults
 - 5,970 hatchery steelhead adults
 - Average of 92% hatchery origin

Overall, the programs are increasing the abundance of steelhead in the ESU


A stylized, layered mountain range graphic in shades of teal and blue, located in the bottom right corner of the slide.

Wenatchee Basin Escapement



Dam count data

Above Wells Dam Escapement




Dam count data


Productivity

- ◆ Methow basin natural-origin steelhead are likely only one generation removed from the hatchery because, on average about 90% of spawners are hatchery origin and it is uncertain what effect this has on productivity
- ◆ Hatchery origin steelhead in excess of available spawning habitat and broodstock needs may decrease productivity

2001 Wenatchee Steelhead Spaw n Timing



Wells Steelhead Spawn Timing 2004




Spatial Structure

- ◆ Radio tag studies indicate program steelhead stray from release areas
- ◆ Studies are examining the effects of release site and/or acclimation periods on the distribution of returning adults
- ◆ Program steelhead have been found spawning in small tributaries
- ◆ Overall, the effect is neutral



Diversity

- ◆ Recent (1996) separation of Wenatchee basin programs from the Wells program is expected to increase diversity over time
 - ◆ High proportion of steelhead on spawning grounds from the Wells program likely decreases local adaptation of the Methow basin and Okanogan basin populations
 - ◆ The small Omak Creek program should increase diversity over time
- 
- A stylized, layered mountain range graphic in shades of teal and blue, located in the bottom right corner of the slide.

Effect by Program

Viability Criteria	Population		
	Wenatchee	Methow	Okanogan
Abundance	↑	↑	↑
Productivity	↔	↔	↔
Spatial Structure	↑	↔	↔
Diversity	↑	↓	↓

Effects of Artificial Propagation on VSP Attributes

Viability Criteria	BRT VSP Risk Score	Decreases Risk	Neutral or Uncertain	Increases Risk
Abundance	3.5	√		
Productivity	4.3		√ ?	√ ?
Spatial Structure	3.1	√		
Diversity	3.6		√	

Endangered Threatened Not Warranted

BRT Findings: 54% 44% 2%

Recommendation: **Change to Threatened**